

AviPro Megan® Vac 1

Indication



For use in broiler, layer and breeder chickens as an immunological aid in the reduction of *Salmonella* Typhimurium, *Salmonella* Enteritidis and *Salmonella* Heidelberg colonisation of the digestive tract and internal organs. Also as an immunological aid in the reduction of *Salmonella* Typhimurium and *Salmonella* Enteritidis colonisation of intestinal, visceral and reproductive tract, including egg colonisation.

Presentation

SV - 500 doses (breeders), 1000 doses (commercial layers) live attenuated *Salmonella* Typhimurium bacteria per vial.
 MV - 1000 doses (breeders), 2000 doses (commercial layers) live attenuated *Salmonella* Typhimurium bacteria per vial.

Vaccination program

Three vaccinations with live attenuated live *Salmonella* Typhimurium during rearing.



Vaccine administration

1 day of age - spray, 2 weeks of age - drinking water or spray, 16 weeks of age - drinking water or spray.

Precautions

Store refrigerated at 2 to 8°C. Do not allow to freeze. Do not expose to disinfectants or heat.

In accordance with the World Organization for Animal Health (OIE) best practice, all susceptible birds on site should be vaccinated concurrently with the dose recommended by the manufacturer.

Vaccinate only healthy birds.

Avoid direct contact with the product. If accidental human contact occurs, clean the area with soap and flush with clean water. Consult a doctor. Wear a mask and gloves while handling the vaccine. Avoid breathing the spray.

See AviPro Megan® Vac 1 package insert for detailed Directions for Use.

AviPro Megan® Vac 1 registered pursuant to the ACVM Act 1997, No. A007935.

See www.foodsafety.govt.nz for registration conditions.

Further Information

Disease Information Sheet

- *Salmonella* Typhimurium

Administration Information Sheet

- Drinking water administration
- Spray administration
- Spray administration of vaccine to day old chicks

