

AviPro Thymovac®

Indication

To induce the production of antibodies to Chicken Anaemia Virus (CAV) by breeder hens and transfer of passive immunity to their progeny.

Presentation



1000 doses live freeze dried Chicken Anemia Virus strain Cux-1 per vial.

Vaccination Program

Layer Breeders and Broiler Breeders receive AviPro Thymovac® vaccine once during rearing.

Vaccine Administration

From 12 until 16 weeks-of-age, but at least 6 weeks prior to the onset of lay administer AviPro Thymovac® in drinking water.



Precautions

Store refrigerated at 2 to 7°C. Do not allow to freeze. Do not expose to disinfectants or heat.

The vaccine virus may spread from vaccinated birds.

Do not administer live CAV vaccine to breeders within 6 weeks of onset of lay. The CAV virus may be transmitted vertically in eggs following vaccination.

Birds less than 3 weeks-of-age infected with live CAV vaccine may display clinical disease, if they lack protection from passive transfer of maternal antibody.

See AviPro Thymovac® package insert for detailed Directions for Use.

AviPro Thymovac® registered pursuant to the ACVM Act 1997, No. A9132. See www.foodsafety.govt.nz for registration conditions.

Further Information

Disease Information Sheet

- Chicken Anaemia Virus (CAV)

Administration Information Sheet

- Drinking water administration

